Post Operative Instructions for Extractions

Joseph J. Havrilla, D.D.S. 905 W. Sproul Rd, Suite #108 Springfield, PA19064 Periodontics and Dental Implants

- 1. Avoid rinsing for the rest of the day. You may begin rinsing **gently** within 24 hours.
- 2. **NO** smoking. **DO NOT** use a straw.
- 3. Strenuous exercise should be avoided until the sutures are removed.
- 4. Chew food on the **opposite** side of the surgical site. Avoid hard and/or crunchy foods.
- 5. Occasional swelling can occur. Apply ice packs as needed at 15-20 minute intervals for the first 48 hours.
- 6. If an antibiotic is prescribed, take according to the directions on the bottle until all of the capsules (tablets) are finished.
- 7. Analgesics for post-operative discomfort are recommended for the first48 hours (can be taken longer)

<u>Ibuprofen</u> (example: Advil, Motrin) 400mg <u>Acetaminophen</u> (ex. Tylenol) 500mg ***Can be taken at the same time (every 4-6 hrs)***

- 8. You may be prescribed additional pain medication by Dr. Havrilla
- 9. Rinse gently with the prescribed mouthwash twice/day. Use the mouthwash until you return for a post-op check. Continue your usual oral hygiene with the rest of your teeth.
- 10.Slight bleeding is not unusual, if necessary, you may apply clean gauze or a wet, warm tea bag, with light even pressure to the site. Call the office if bleeding continues.
- 11. If you have any questions or concerns, you may call the office at (610)328-9608 or Doctor can be reached on his cell phone at (484)716-3403, he will return your call as soon as possible.

***We encourage our patients to contact us with post operative concerns.

PLEASE DO NOT WAIT!

Copies of our instructions are also on our website www.JJHPerio.com

Post Operative Instructions for Extractions